

## LIST OF TABLES

Table III.1. Total number of fish caught by gill net during spring or fall season; number of fish measured; and number of fish in panels where none were measured. . . . .	15
Table III.2. Species, size ranges, months or season, and gears from which data were selected for trend analysis. . . . .	18
Table III.3. Summary of CPUE definitions used for analysis of species caught by gill net. . . . .	20
Table III.4. Summary of estimated values of l and k used in analyses. . . . .	22
Table III.5. Analysis of deviance for young-of-the-year blue crab caught by bag seine, January-December, 1978-1990. . . . .	35
Table III.6. Analysis of deviance for young-of-the-year blue crab caught by trawl, January-December, 1983-1990. . . . .	35
Table III.7. Analysis of deviance for juvenile blue crab caught by trawl, January-December, 1983-1990. . . . .	36
Table III.8. Analysis of deviance for first-time spawning blue crab caught by trawl, January-December, 1983-1990. . . . .	36
Table III.9. Analysis of deviance for first-time spawning blue crab caught by gill net, spring season, 1983-1990. . . . .	38
Table III.10. Analysis of deviance for remaining adult blue crab caught by gill net, spring season, 1983-1990. . . . .	38
Table III.11. Analysis of deviance for brown shrimp, all size classes, caught by trawl, April to November, 1963-1968. . . . .	54
Table III.12. Analysis of deviance for brown shrimp, all size classes, caught by trawl, April to November, 1972-1980. . . . .	54
Table III.13. Analysis of deviance for young-of-the-year brown shrimp caught by bag seine, April-November, 1978-1990. . . . .	55
Table III.14. Analysis of deviance for first-time spawning brown shrimp caught by trawl, May to August, 1963-1968. . . . .	55
Table III.15. Analysis of deviance for first-time spawning brown shrimp caught by trawl, May to August, 1972-1980. . . . .	56

Table III.16. Analysis of deviance for first-time spawning brown shrimp caught by trawl, May-August, 1983-1990. . . . .	56
Table III.17. Analysis of deviance for adult grass shrimp caught by bag seine, January-December, 1983-1990. . . . .	64
Table III.18. Analysis of deviance for white shrimp, all size classes, caught by trawl, June to November, 1963-1968. . . . .	77
Table III.19. Analysis of deviance for white shrimp, all size classes, caught by trawl, June to November, 1972-1980. . . . .	77
Table III.20. Analysis of deviance for young-of-the-year white shrimp caught by bag seine, June-December, 1977-1990. . . . .	78
Table III.21. Analysis of deviance for juvenile white shrimp caught by trawl, July-April, 1982-1990. . . . .	78
Table III.22. Analysis of deviance for first-time spawning white shrimp caught by trawl, August to November, 1963-1968. . . . .	79
Table III.23. Analysis of deviance for first-time spawning white shrimp caught by trawl, August to November, 1972-1980. . . . .	79
Table III.24. Analysis of deviance for first-time spawning white shrimp caught by trawl, April-May, 1982-1990. . . . .	80
Table III.25. Analysis of deviance for first-time spawning white shrimp caught by trawl, August-November, 1982-1990. . . . .	80
Table III.26. Analysis of deviance for young-of-the-year Atlantic croaker caught by bag seine, December-March, 1977-1989. . . . .	94
Table III.27. Analysis of deviance for first-time spawning Atlantic croaker caught by trawl, March-September, 1983-1990. . . . .	94
Table III.28. Analysis of deviance for remaining adult Atlantic croaker caught by gill net, 230-275 mm, fall season, 1975-1989. . . . .	95
Table III.29. Analysis of deviance for juvenile bay anchovy caught by bag seine, May-October, 1978-1990. . . . .	108
Table III.30. Analysis of deviance for juvenile bay anchovy caught by trawl, December-March, 1983-1990. . . . .	108

Table III.31. Analysis of deviance for first-time spawning bay anchovy caught by bag seine, April-October, 1978-1990. . . . .	109
Table III.32. Analysis of deviance for first-spawning bay anchovy caught by trawl, January-December, 1983-1990. . . . .	109
Table III.33. Analysis of deviance for remaining adult bay anchovy caught by bag seine, April-November, 1978-1990. . . . .	110
Table III.34. Analysis of deviance for remaining adult bay anchovy caught by trawl, January-December, 1983-1990. . . . .	110
Table III.35. Analysis of deviance for juvenile black drum caught by bag seine, June-August, 1978-1990. . . . .	124
Table III.36. Analysis of deviance for first-time spawning black drum caught by gill net, fall season, 1975-1989. . . . .	124
Table III.37. Analysis of deviance for remaining adult black drum caught by gill net, fall season, 1975-1989. . . . .	125
Table III.38. Analysis of deviance for young-of-the-year Gulf menhaden caught by bag seine, February-July, 1978-1989. . . . .	135
Table III.39. Analysis of deviance for first-time spawning Gulf menhaden caught by trawl, January-December, 1983-1990. . . . .	135
Table III.40. Analysis of deviance for remaining adult Gulf menhaden caught by gill net, fall season, 1975-1989. . . . .	136
Table III.41. Analysis of deviance for young-of-the-year pinfish caught by bag seine, March-June, 1978-1989. . . . .	149
Table III.42. Analysis of deviance for juvenile pinfish caught by trawl, June-August, 1983-1990. . . . .	149
Table III.43. Analysis of deviance for first-time spawning pinfish caught by trawl, September-November, 1983-1990. . . . .	150
Table III.44. Analysis of deviance for young-of-the-year red drum caught by bag seine, October-January, 1977-1989. . . . .	155
Table III.45. Analysis of deviance for juvenile red drum caught by gill net, spring season, 1983-1990. . . . .	156
Table III.46. Analysis of deviance for first-time spawning red drum caught by gill net, spring season, 1983-1990. . . . .	156

Table III.47. Analysis of deviance for young-of-the-year sand seatrout caught by bag seine, April-October, 1978-1989. . . . .	168
Table III.48. Analysis of deviance for juvenile sand seatrout caught by trawl, May-December, 1983-1990. . . . .	168
Table III.49. Analysis of deviance for first-time spawning sand seatrout caught by trawl, April-November, 1983-1990. . . . .	169
Table III.50. Analysis of deviance for remaining adult sand seatrout caught by gill net, fall season, 1975-1990. . . . .	169
Table III.51. Analysis of deviance for young-of-the-year southern flounder caught by bag seine, February-March, 1978-1989. . . . .	179
Table III.52. Analysis of deviance table for adult southern flounder caught by gill net, fall season, 1975-1989. . . . .	179
Table III.53. Analysis of deviance for young-of-the-year spotted seatrout caught by bag seine, June-November, 1978-1990. . . . .	187
Table III.54. Analysis of deviance for first-time spawning spotted seatrout caught by gill net, spring season, 1977-1990. . . . .	188
Table III.55. Analysis of deviance for remaining adult spotted seatrout caught by gill net, spring season, 1977-1990. . . . .	188
Table III.56. Analysis of deviance for young-of-the-year striped mullet caught by bag seine, February-March, 1978-1990. . . . .	199
Table III.57. Analysis of deviance for first-time spawning striped mullet caught by trawl, December-March, 1983-1990. . . . .	199
Table III.58. Analysis of deviance for Age IV striped mullet caught by gill net, fall season, 1975-1989. . . . .	201
Table III.59. Analysis of deviance for Age V+ striped mullet caught by gill net, fall season, 1975-1989. . . . .	201
Table III.60. Commercial landings reported from the Galveston Estuary during 1890 and 1989. . . . .	208
Table IV.1. Analysis of variance for black-bellied plovers. . . . .	283
Table IV.2. Analysis of variance for willets. . . . .	283
Table IV.3. Analysis of variance for sanderlings. . . . .	284

Table IV.4. Analysis of variance for western sandpipers. . . . .	284
Table IV.5. Analysis of variance for American avocets. . . . .	285
Table IV.6. Analysis of variance for dunlins. . . . .	285
Table IV.7. Analysis of variance for tricolored herons. . . . .	292
Table IV.8. Analysis of variance for snowy egrets. . . . .	293
Table IV.9. Analysis of variance for roseate spoonbills. . . . .	293
Table IV.10. Analysis of variance for black skimmers. . . . .	296
Table IV.11. Analysis of variance for olivaceous cormorants. . . . .	297
Table IV.12. Analysis of variance for black-crowned night-herons. . . . .	297
Table IV.13. Analysis of variance for Forster's terns. . . . .	301
Table IV.14. Analysis of variance for great egrets. . . . .	301
Table IV.15. Analysis of variance for merganser spp. . . . .	308
Table IV.16. Analysis of variance for American coots. . . . .	315
Table IV.17. Analysis of variance for green-winged teal. . . . .	315
Table IV.18. Analysis of variance for northern shovelers. . . . .	316
Table IV.19. Analysis of variance for American wigeons. . . . .	316
Table IV.20. Analysis of variance for mallards. . . . .	318
Table IV.21. Analysis of variance for gadwalls. . . . .	318
Table IV.22. Analysis of variance for mottled ducks. . . . .	320
Table IV.23. Analysis of variance for northern pintails. . . . .	326
Table IV.24. Analysis of variance for blue-winged teal. . . . .	326
Table IV.25. Analysis of variance for scaup spp. . . . .	328
Table IV.26. Analysis of variance for ruddy ducks. . . . .	328
Table IV.27. Analysis of variance for buffleheads. . . . .	328

Table IV.28. Analysis of variance for wood ducks. . . . .	333
Table IV.29. Analysis of variance for ring-necked ducks. . . . .	333
Table IV.30. Analysis of variance for canvasbacks. . . . .	334
Table IV.31. Analysis of variance for fulvous whistling ducks. . . . .	334
Table IV.32. Analysis of variance for red-winged blackbirds. . . . .	334
Table IV.33. Analysis of variance for American alligators. . . . .	337
Table V.1. Dominant genera of phytoplankton in Galveston Bay in 1969. . . . .	351
Table V.2. Surface primary production rates determined at each sampling station from October 1972 to April 1974. . . . .	358
Table V.3. A list of zooplankton species collected in the Galveston Estuary during the period June 1974 through September 1975. . . . .	361
Table V.4. A comparison of phytoplankton assemblages, cell concentrations and primary production rates for various Texas estuaries. . . . .	362
Table V.5. Annual phytoplankton primary production estimates for selected North American estuaries. . . . .	363
Table V.6. Comparison of zooplankton biomass and abundances for several Texas estuaries.	
A) Comparison of data from UTMSI TWDB studies over the past four years. . . . .	365
B) Comparison of other Texas estuary studies. . . . .	366
C) Comparison of <i>Acartia tonsa</i> abundance in Texas estuaries. . . . .	366
Table V.7. Comparison of zooplankton abundances for several U.S. estuaries. . . . .	367
Table VI.1. Abundances of infauna in sediment cores from marsh sites in the Galveston Estuary. . . . .	381
Table VI.2. Annual variation in percent abundance of annelid worms among infauna of three types of habitats associated with a salt marsh on Galveston Island. . . . .	388
Table VI.3. Annual variation in percent abundance of peracarid crustaceans among infauna of three types of habitats associated with a salt marsh on Galveston Island. . . . .	389

Table VI.4. Annual variation in percent abundance of mollusks among infauna of three types of habitats associated with a salt marsh on Galveston Island. . . . .	390
Table VII.1. Macrobenthos studies conducted in the Galveston Estuary and adjacent bay systems. . . . .	416
Table VII.2. Frequency of sampling at Texas Water Commission stations within the Galveston Estuary and the Brazos River. . . . .	445