

LIST OF FIGURES

Figure III.1. A. Mean annual CPUE and predicted CPUE for young-of-the-year blue crab caught by bag seine. B. Mean CPUE by month.	27
Figure III.2. A. Mean annual CPUE and predicted CPUE for young-of-the-year blue crab caught by trawl. B. Mean CPUE by month.	28
Figure III.3. A. Mean annual CPUE and predicted CPUE for juvenile blue crab caught by trawl. B. Mean CPUE by month.	29
Figure III.4. A. Mean annual CPUE and predicted CPUE for first-time spawning blue crab caught by trawl. B. Mean CPUE by month.	30
Figure III.5. A. Mean seasonal (spring) and predicted CPUE for first-time spawning blue crab caught by gill net. B. Mean catch per hour gill net set duration.	31
Figure III.6. A. Mean seasonal (spring) and predicted CPUE for remaining adult blue crab caught by gill net. B. Mean catch per hour gill net set duration.	32
Figure III.7. Mean CPUE by month for four size classes of blue crab caught by bag seine.	33
Figure III.8. Mean CPUE by month for four size classes of blue crab caught by trawl.	33
Figure III.9. Size frequency distribution for blue crab caught by gill net. . .	34
Figure III.10. Distribution of young-of-the-year blue crab caught by bag seine.	39
Figure III.11. Distribution of young-of-the-year blue crab caught by trawl. .	40
Figure III.12. Distribution of juvenile blue crab caught by trawl.	41
Figure III.13. Distribution of first-time spawning blue crab caught by trawl.	42
Figure III.14. Distribution of first-time spawning blue crab caught by gill net.	43
Figure III.15. Distribution of remaining adult blue crab caught by gill net.	44

Figure III.16. A. Mean seasonal (April-November) CPUE and predicted CPUE for all sizes of brown shrimp caught by trawl, 1963-1968. B. Mean CPUE by month.	46
Figure III.17. A. Mean seasonal (May-August) CPUE and predicted CPUE for first-time spawning brown shrimp caught by trawl, 1963-1968. B. Mean CPUE by month.	47
Figure III.18. A. Mean seasonal (April-November) CPUE and predicted CPUE for all sizes of brown shrimp caught by trawl, 1972-1980. B. Mean CPUE by month.	48
Figure III.19. A. Mean seasonal (May-August) CPUE and predicted CPUE for first-time spawning brown shrimp caught by trawl, 1972-1980. B. Mean CPUE by month.	49
Figure III.20. A. Mean seasonal (April-November) CPUE and predicted CPUE for young-of-the-year brown shrimp caught by bag seine. B. Mean CPUE by month.	50
Figure III.21. A. Mean seasonal (May-August) CPUE and predicted CPUE for first-time spawning brown shrimp caught by trawl, 1983-1990. B. Mean CPUE by month.	51
Figure III.22. Mean CPUE by month for three size classes of brown shrimp caught by bag seine.	53
Figure III.23. Mean CPUE by month for four size classes of brown shrimp caught by trawl.	53
Figure III.24. Distribution of young-of-the-year brown shrimp caught by bag seine.	58
Figure III.25. Distribution of first-time spawning brown shrimp caught by trawl.	59
Figure III.26. A. Mean seasonal CPUE exclusive of zero catches, percentage zero catches, and predicted values for young-of-the-year brown shrimp caught by bag seine. B. Mean seasonal CPUE exclusive of zero catches, percentage zero catches, and predicted values for first-time spawning brown shrimp caught by trawl.	61
Figure III.27. A. Mean annual CPUE and predicted CPUE for adult grass shrimp caught by bag seine. B. Mean CPUE by month.	63
Figure III.28. Mean CPUE by month for three size classes of grass shrimp caught by bag seine.	64

Figure III.29. Distribution of adult grass shrimp caught by bag seine.	65
Figure III.30. A. Mean seasonal (June-November) CPUE and predicted CPUE for all sizes of white shrimp caught by trawl, 1963-1968. B. Mean CPUE by month.	67
Figure III.31. A. Mean seasonal (August-November) CPUE and predicted CPUE for first-time spawning white shrimp caught by trawl, 1963-1968. B. Mean CPUE by month.	68
Figure III.32. A. Mean seasonal (June-November) and predicted CPUE for all sizes of white shrimp caught by trawl, 1972-1980. B. Mean CPUE by month.	69
Figure III.33. A. Mean seasonal (August-November) CPUE and predicted CPUE for first-time spawning white shrimp caught by trawl, 1972-1980. B. Mean CPUE by month.	70
Figure III.34. A. Mean seasonal (June-December) CPUE and predicted CPUE for young-of-the-year white shrimp caught by bag seine. B. Mean CPUE by month.	71
Figure III.35. A. Mean seasonal (July-April) CPUE and predicted CPUE for juvenile white shrimp caught by trawl, 1982-1990. B. Mean CPUE by month.	72
Figure III.36. A. Mean seasonal (April-May) CPUE and predicted CPUE for first-time spawning white shrimp caught by trawl, 1982-1990. B. Mean CPUE by month.	73
Figure III.37. Mean seasonal (August-November) and predicted CPUE for first-time spawning white shrimp caught by trawl, 1982-1990.	74
Figure III.38. Mean CPUE by month for four size classes of white shrimp caught by bag seine.	75
Figure III.39. Mean CPUE by month for four size classes of white shrimp caught by trawl.	75
Figure III.40. Distribution of young-of-the-year white shrimp caught by bag seine.	81
Figure III.41. Distribution of juvenile white shrimp caught by trawl.	82
Figure III.42. Distribution of first-time spawning white shrimp caught by trawl, April-May.	83

Figure III.43. Distribution of first-time spawning white shrimp caught by trawl, August-November.	84
Figure III.44. A. Mean seasonal CPUE exclusive of zero catches, percentage zero catches, and predicted values for young-of-the-year white shrimp caught by bag seine. B. Mean seasonal CPUE exclusive of zero catches, percentage zero catches, and predicted values for juvenile white shrimp caught by trawl.	86
Figure III.45. A. Mean seasonal (April-May) CPUE exclusive of zero catches, percentage zero catches, and predicted values for first-time spawning white shrimp caught by trawl. B. Mean seasonal (August-November) CPUE exclusive of zero catches, percentage zero catches, and predicted values for first-time spawning white shrimp caught by trawl.	87
Figure III.46. A. Mean seasonal (December-March) CPUE and predicted CPUE for young-of-the-year Atlantic croaker caught by bag seine. B. Mean CPUE by month.	89
Figure III.47. A. Mean seasonal (March-September) CPUE and predicted CPUE with confidence intervals for first-time spawning Atlantic croaker caught by trawl. B. Mean CPUE by month.	90
Figure III.48. A. Mean seasonal (fall) CPUE and predicted CPUE for remaining adult Atlantic croaker caught by gill net. B. Mean catch per hour gill net set duration.	91
Figure III.49. Mean CPUE by month for five size classes of Atlantic croaker caught by bag seine.	92
Figure III.50. Mean CPUE by month for four size classes of Atlantic croaker caught by trawl.	92
Figure III.51. Size frequency distribution for Atlantic croaker caught by gill net.	93
Figure III.52. Distribution of young-of-the-year Atlantic croaker caught by bag seine.	96
Figure III.53. Distribution of first-time spawning Atlantic croaker caught by trawl.	97
Figure III.54. Distribution of remaining adult Atlantic croaker caught by gill net.	98

Figure III.55. A. Mean seasonal (May-October) CPUE and predicted CPUE for juvenile bay anchovy caught by bag seine. B. Mean CPUE by month	100
Figure III.56. A. Mean seasonal (April-October) CPUE and predicted CPUE for first-time spawning bay anchovy caught by bag seine. B. Mean CPUE by month.	101
Figure III.57. A. Mean seasonal (April-November) CPUE and predicted CPUE for remaining adult bay anchovy caught by bag seine. B. Mean CPUE by month.	102
Figure III.58. A. Mean seasonal (December-March) and fitted CPUE for juvenile bay anchovy caught by trawl. B. Mean CPUE by month.	103
Figure III.59. A. Mean annual CPUE and predicted CPUE for first-time spawning bay anchovy caught by trawl. B. Mean CPUE by month.	104
Figure III.60. A. Mean annual CPUE and predicted CPUE for remaining adult bay anchovy caught by trawl. B. Mean CPUE by month.	105
Figure III.61. Mean CPUE by month for three sizes of bay anchovy caught by bag seine.	106
Figure III.62. Mean CPUE by month for three sizes of bay anchovy caught by trawl.	106
Figure III.63. Distribution of juvenile bay anchovy caught by bag seine.	112
Figure III.64. Distribution of first-time spawning bay anchovy caught by bag seine.	113
Figure III.65. Distribution of remaining adult bay anchovy caught by bag seine.	114
Figure III.66. Distribution of young-of-the-year bay anchovy caught by trawl.	115
Figure III.67. Distribution of first-time spawning bay anchovy caught by trawl.	116
Figure III.68. Distribution of remaining adult bay anchovy caught by trawl.	117

Figure III.69. A. Mean seasonal (June-August) CPUE and predicted annual CPUE for young-of-the-year black drum caught by bag seine. B. Mean CPUE by month.	119
Figure III.70. A. Mean seasonal (fall) CPUE and predicted CPUE for first-time spawning black drum caught by gill net. B. Mean catch per hour gill net set duration.	120
Figure III.71. A. Mean seasonal (fall) CPUE and predicted CPUE for remaining adult black drum caught by gill net. B. Mean catch per hour gill net set duration.	121
Figure III.72. Mean CPUE by month for three size classes of black drum caught by bag seine.	122
Figure III.73. Size frequency distribution for black drum caught by gill net during fall sampling period.	122
Figure III.74. Distribution of young-of-the-year black drum caught by bag seine.	126
Figure III.75. Distribution of first-time spawning black drum caught by gill net.	127
Figure III.76. Distribution of remaining adult black drum caught by gill net.	128
Figure III.77. A. Mean seasonal (February-July) CPUE and predicted CPUE for young-of-the-year Gulf menhaden caught by bag seine. B. Mean CPUE by month.	130
Figure III.78. A. Mean annual CPUE and predicted CPUE for first-time spawning Gulf menhaden caught by trawl. B. Mean CPUE by month.	131
Figure III.79. A. Mean seasonal (fall) CPUE and predicted CPUE for remaining adult Gulf menhaden (all sizes) caught by gill net. B. Mean catch per hour gill net set duration.	132
Figure III.80. Mean CPUE by month for five size classes of Gulf menhaden caught by bag seine.	133
Figure III.81. Mean CPUE by month for four size classes of Gulf menhaden caught by trawl.	133
Figure III.82. Size frequency distribution of Gulf menhaden caught by gill net.	134

Figure III.83. Distribution of young-of-the-year Gulf menhaden caught by bag seine.	137
Figure III.84. Distribution of first-time spawning Gulf menhaden caught by trawl.	138
Figure III.85. Distribution of remaining adult Gulf menhaden caught by gill net.	139
Figure III.86. A. Mean seasonal (March-June) CPUE and predicted CPUE for young-of-the-year pinfish caught by bag seine. B. Mean CPUE by month.	141
Figure III.87. A. Mean seasonal (June-August) CPUE and predicted CPUE for juvenile pinfish caught by trawl. B. Mean CPUE by month.	142
Figure III.88. A. Mean seasonal (September-November) CPUE and predicted CPUE for first-time spawning pinfish caught by trawl. B. Mean CPUE by month.	143
Figure III.89. Mean CPUE by month for four size classes of pinfish caught by bag seine.	144
Figure III.90. Mean CPUE by month for four size classes of pinfish caught by trawl.	144
Figure III.91. Distribution of young-of-the-year pinfish caught by bag seine.	146
Figure III.92. Distribution of juvenile pinfish caught by trawl.	147
Figure III.93. Distribution of first-time spawning pinfish caught by trawl.	148
Figure III.94. A. Mean seasonal (October-January) CPUE and predicted CPUE for young-of-the-year red drum caught by bag seine. B. Mean CPUE by month.	151
Figure III.95. A. Mean seasonal (spring) CPUE and predicted CPUE for juvenile red drum caught by gill net. B. Mean catch per hour gill net set duration.	152
Figure III.96. A. Mean seasonal (spring) CPUE and predicted CPUE for first-time spawning red drum caught by gill net. B. Mean catch per hour gill net set duration.	153
Figure III.97. Mean CPUE by month for three sizes of red drum caught by bag seine.	154

Figure III.98. Size frequency distribution of red drum caught by gill net.	154
Figure III.99. Distribution of young-of-the-year red drum caught by bag seine.	158
Figure III.100. Distribution of juvenile red drum caught by gill net.	159
Figure III.101. Distribution of first-time spawning red drum caught by gill net.	160
Figure III.102. A. Mean seasonal (April-October) CPUE and predicted CPUE for young-of-the-year sand seatrout caught by bag seine. B. Mean CPUE by month.	162
Figure III.103. A. Mean seasonal (May-December) CPUE and predicted CPUE for juvenile sand seatrout caught by trawl. B. Mean CPUE by month.	163
Figure III.104. A. Mean seasonal (April-November) CPUE and predicted CPUE for first-time spawning sand seatrout caught by trawl. B. Mean CPUE by month.	164
Figure III.105. A. Mean seasonal (fall) CPUE and predicted CPUE for remaining adult sand seatrout caught by gill net. B. Mean catch per hour gill net set duration.	165
Figure III.106. Mean CPUE by month for three size classes of sand seatrout caught by bag seine.	166
Figure III.107. Mean CPUE by month for four size classes of sand seatrout caught by trawl.	166
Figure III.108. Size frequency distribution for sand seatrout caught by gill net.	167
Figure III.109. Distribution of young-of-the-year sand seatrout caught by bag seine.	171
Figure III.110. Distribution of juvenile sand seatrout caught by trawl.	172
Figure III.111. Distribution of first-time spawning sand seatrout caught by trawl.	173
Figure III.112. Distribution of remaining adult sand seatrout caught by gill net.	174
Figure III.113. Mean length for sand seatrout caught by trawl.	170

Figure III.114. A. Mean seasonal (February-March) CPUE and predicted CPUE for young-of-the-year southern flounder caught by bag seine. B. Mean CPUE by month.	176
Figure III.115. A. Mean seasonal (fall) CPUE and predicted CPUE for adult southern flounder caught by gill net. B. Mean catch per hour gill net set duration.	177
Figure III.116. Mean CPUE by month for three size classes of southern flounder caught by bag seine.	178
Figure III.117. Size frequency distribution for southern flounder caught by gill net.	178
Figure III.118. Distribution of young-of-the-year southern flounder caught by bag seine.	181
Figure III.119. Distribution of adult southern flounder caught by gill net.	182
Figure III.120. A. Mean seasonal (June-November) CPUE and predicted CPUE for young-of-the-year spotted seatrout caught by bag seine. B. Mean CPUE by month.	183
Figure III.121. A. Mean seasonal (spring) CPUE and predicted CPUE for first-time spawning spotted seatrout caught by gill net. B. Mean catch per hour gill net set duration.	184
Figure III.122. A. Mean seasonal (spring) CPUE and predicted CPUE for remaining adult spotted seatrout caught by gill net. B. Mean catch per hour gill net set duration.	185
Figure III.123. Mean CPUE by month for four size classes of spotted seatrout caught by bag seine.	186
Figure III.124. Size frequency distribution of spotted seatrout caught by gill net.	186
Figure III.125. Distribution of young-of-the-year spotted seatrout caught by bag seine.	190
Figure III.126. Distribution of first-time spawning spotted seatrout caught by gill net.	191
Figure III.127. Distribution of remaining adult spotted seatrout caught by gill net.	192

Figure III.128. A. Mean seasonal (February-March) CPUE and predicted CPUE for young-of-the-year striped mullet caught by bag seine. B. Mean CPUE by month.	193
Figure III.129. A. Mean seasonal (December-March) CPUE and predicted CPUE for first-time spawning striped mullet caught by trawl. B. Mean CPUE by month.	194
Figure III.130. A. Mean seasonal (fall) CPUE and predicted CPUE for Age IV adult striped mullet caught by gill net. B. Mean catch per hour gill net set duration.	195
Figure III.131. A. Mean seasonal (fall) CPUE and predicted CPUE for Age V+ adult striped mullet caught by gill net. B. Mean catch per hour gill net set duration.	196
Figure III.132. Mean CPUE by month for four size classes of striped mullet caught by bag seine.	197
Figure III.133. Mean CPUE by month for four size classes of striped mullet caught by trawl.	197
Figure III.134. Size frequency distribution for striped mullet caught by gill net.	198
Figure III.135. Distribution of young-of-the-year striped mullet caught by bag seine.	202
Figure III.136. Distribution of first-time spawning striped mullet caught by trawl.	203
Figure III.137. Distribution of Age IV adult striped mullet caught by gill net.	204
Figure III.138. Distribution of Age V+ adult striped mullet caught by gill net.	205
Figure III.3.1. Species composition, by weight, of commercial landings. . .	235
Figure III.3.2. Commercial landings of white shrimp and brown and pink shrimp.	236
Figure III.3.3. Commercial landings of white shrimp and brown and pink shrimp in the Gulf of Mexico offshore of the Galveston Estuary.	236
Figure III.3.4. Commercial landings of American oysters.	237

Figure III.3.5. Commercial landings of blue crab.	237
Figure III.3.6. Commercial landings of squid.	239
Figure III.3.7. Commercial landings of stone crab.	239
Figure III.3.8. Commercial landings of all finfish.	240
Figure III.3.9. Species composition of commercial finfish landings.	240
Figure III.3.10. Commercial landings of spotted and sand seatrout.	241
Figure III.3.11. Commercial landings of red drum.	241
Figure III.3.12. Commercial landings of black drum.	242
Figure III.3.13. Commercial landings of flounder.	242
Figure III.3.14. Commercial landings of sheepshead.	243
Figure III.3.15. Commercial landings of Atlantic croaker.	243
Figure III.3.16. Commercial landings of kingfish (whiting).	244
Figure III.3.17. Commercial landings of unclassified food and scrap.	244
Figure III.3.18. Commercial landings of spotted and sand seatrout.	245
Figure III.3.19. Commercial landings of mullet.	245
Figure III.3.20. Commercial landings of gafftopsail catfish.	246
Figure III.3.21. Recreational landings of all finfish caught by private sport-boat fishermen.	246
Figure III.3.22. Recreational effort expended by private sport-boat fishermen.	247
Figure III.3.23. Species composition of recreational finfish landings caught by private sport-boat fishermen.	247
Figure III.3.24. Recreational landings of Atlantic croaker caught by private sport-boat fishermen.	248
Figure III.3.25. Recreational landings of sand seatrout caught by private sport-boat fishermen.	248

Figure III.3.26. Recreational landings of spotted seatrout caught by private sport-boat fishermen.	249
Figure III.3.27. Recreational landings of black drum caught by private sport-boat fishermen.	249
Figure III.3.28. Recreational landings of sheepshead caught by private sport-boat fishermen.	250
Figure III.3.29. Recreational landings of gafftopsail catfish caught by private sport-boat fishermen.	250
Figure III.3.30. Recreational landings of southern flounder caught by private sport-boat fishermen.	251
Figure III.3.31. Recreational landings of red drum caught by private sport-boat fishermen.	251
Figure IV.1. Data set coverage of birds and alligators in the Galveston Estuary.	266
Figure IV.2. A. Number of black-bellied plovers per survey during Shorebird Surveys of Bolivar Flats. B. Trend in black-bellied plovers from 1980 to 1990 during Shorebird Surveys of Bolivar Flats.	272
Figure IV.3. A. The number of black-bellied plovers per count during Christmas Bird Counts. B. Trend in black-bellied plovers from 1965 to 1989 during Christmas Bird Counts.	273
Figure IV.4. A. Number of willets per survey during Shorebird Surveys of Bolivar Flats. B. Trend in willets from 1980 to 1990 during Shorebird Surveys of Bolivar Flats.	274
Figure IV.5. A. The number of willets per count during Christmas Bird Counts. B. Trend in willets from 1965 to 1989 during Christmas Bird Counts.	275
Figure IV.6. A. Number of sanderlings per survey during Shorebird Surveys of Bolivar Flats. B. Trend in sanderlings from 1980 to 1990 during Shorebird Survey of Bolivar Flats.	276
Figure IV.7. A. The number of sanderlings per count during Christmas Bird Counts. B. Trend in sanderlings from 1965 to 1989 during Christmas Bird Counts.	277

Figure IV.8. A. Number of western sandpipers per survey during Shorebird Surveys of Bolivar Flats. B. Trend of western sandpipers from 1980 to 1990 during Shorebird Surveys of Bolivar Flats.	278
Figure IV.9. A. Number of western sandpipers per count during Christmas Bird Counts. B. Trend in western sandpipers from 1965 to 1989 during Christmas Bird Counts.	279
Figure IV.10. A. Individual counts and mean number of American avocets per survey during Shorebird Surveys of Bolivar Flats. B. Individual counts and mean number of dunlins per survey during Shorebird Surveys of Bolivar Flats.	280
Figure IV.11. A. The number of American avocets per count during Christmas Bird Counts. B. Trend in American avocets from 1965 to 1989 during Christmas Bird Counts.	281
Figure IV.12. A. The number of dunlins per count during Christmas Bird Counts. B. Trend in dunlins from 1965 to 1989 during Christmas Bird Counts.	282
Figure IV.13. A. Trend in tricolored herons per year and the trend in numbers of colonies containing tricolored herons from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and trend in tricolored herons per colony from 1973 to 1990 during Texas Colonial Waterbird Surveys.	286
Figure IV.14. A. Individual counts and mean number of black-crowned night herons per count during Christmas Bird Counts. B. Individual counts and mean number of tricolored herons per count during Christmas Bird Counts.	287
Figure IV.15. A. Trend in snowy egrets per year and the number of colonies containing snowy egrets from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and trend in snowy egrets per colony from 1973 to 1990 during Texas Colonial Waterbird Surveys.	288
Figure IV.16. A. The number of snowy egrets per count during Christmas Bird Counts. B. Trend in snowy egrets from 1965 to 1989 during Christmas Bird Counts.	289
Figure IV.17. A. Total number of roseate spoonbills per year and the number of colonies containing roseate spoonbills during Texas Colonial Waterbird Surveys. B. Individual colony counts and trend in roseate spoonbills per colony from 1973 to 1990 during Texas Colonial Waterbird Surveys.	290

Figure IV.18. A. The number of roseate spoonbills per count during Christmas Bird Counts. B. Trend in roseate spoonbills from 1965 to 1989 during Christmas Bird Counts.	291
Figure IV.19. A. Trend in numbers of black skimmers per year and the number of colonies containing black skimmers from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and trend in numbers of black skimmers per colony from 1973 to 1990 during Texas Colonial Waterbird Surveys.	294
Figure IV.20. A. Individual counts and mean number of black skimmers per count during Christmas Bird Counts. B. Individual counts and mean number of fulvous whistling ducks per count during Christmas Bird Counts.	295
Figure IV.21. A. Trend in olivaceous cormorants per year and the trend in number of colonies containing olivaceous cormorants from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and mean number of olivaceous cormorants per colony during Texas Colonial Waterbird Surveys.	298
Figure IV.22. A. The number of olivaceous cormorants per count during Christmas Bird Counts. B. Trend in olivaceous cormorants from 1965 to 1989 during Christmas Bird Counts.	299
Figure IV.23. A. Total number of black-crowned night-herons per year and the trend in number of colonies containing black-crowned night-herons from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and mean number of black-crowned night-herons per colony during Texas Colonial Waterbird Surveys.	300
Figure IV.24. A. Total number of Forster's terns per year and the trend in number of colonies containing Forster's terns from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and mean number of Forster's terns per colony during Texas Colonial Waterbird Surveys.	302
Figure IV.25. A. The number of Forster's terns per count during Christmas Bird Counts. B. Trend in Forster's terns from 1965 to 1989 during Christmas Bird Counts.	303
Figure IV.26. A. Total number of great egrets per year and the trend in number of colonies containing great egrets from 1973 to 1990 during Texas Colonial Waterbird Surveys. B. Individual colony counts and mean number of great egrets per colony during Texas Colonial Waterbird Surveys.	304

Figure IV.27. A. The number of great egrets per count during Christmas Bird Counts. B. Trend in great egrets from 1965 to 1989 during Christmas Bird Counts.	305
Figure IV.28. A. Plot of ordination scores of colonial waterbird species surveyed from 1973 to 1990. B. Plot of ordination scores of the sample years for Texas Waterbird Colonies from 1973 to 1990.	307
Figure IV.29. Individual counts and trend in merganser spp. from 1986 to 1991 during Mid-winter Waterfowl Cruise Counts.	308
Figure IV.30. A. The number of mergansers spp. per count during Christmas Bird Counts. B. Trend in mergansers from 1965 to 1989 during Christmas Bird Counts.	309
Figure IV.31. A. Individual counts and mean number of American wigeons per transect during Med-winter Waterfowl Transects. B. Individual counts and mean number of American coots per transect from 1986 to 1991 during Min-winter Waterfowl Transects.	310
Figure IV.32. A. Individual counts and mean number of northern shovelers per count during Christmas Bird Counts. B. Individual counts and mean number of American coots per count during Christmas Bird Counts.	311
Figure IV.33. A. Individual counts and mean number of green-winged teal per transect during Mid-winter Waterfowl Transects. B. Individual counts and mean number of mallards per transect during Mid-winter Waterfowl Transects.	312
Figure IV.34. A. Individual counts and mean number of green-winged teal per count during Christmas Bird Counts. B. Individual counts and mean number of American wigeons per count during Christmas Bird Counts.	313
Figure IV.35. A. Individual counts and mean number of northern shovelers per transect during Mid-winter Waterfowl Transects. B. Individual counts and mean number of gadwalls per transect during Mid-winter Waterfowl Transects.	314
Figure IV.36. A. The number of mallards per count during Christmas Bird Counts. B. Trend in mallards from 1965 to 1989 during Christmas Bird Counts.	317
Figure IV.37. A. The number of gadwalls per count during Christmas Bird Counts. B. Trend in gadwalls from 1965 to 1989 during Christmas Bird Counts.	319

Figure IV.38. Individual counts and trend in mottled ducks from 1986 to 1991 during Mid-winter Waterfowl Transects.	320
Figure IV.39. A. Number of mottled ducks per count during Christmas Bird Counts. B. Trend in mottled ducks from 1965 to 1989 during Christmas Bird Counts.	321
Figure IV.40. A. The numbers of northern pintails per transect during Mid-winter Waterfowl Transects. B. Trend in northern pintails from 1986 to 1991 during Mid-winter Waterfowl Transects.	322
Figure IV.41. A. The number of northern pintails per count during Christmas Bird Counts. B. Trend in northern pintails from 1965 to 1989 during Christmas Bird Counts.	323
Figure IV.42. A. The numbers of blue-winged teal per transect during Mid-winter Waterfowl Transects. B. Trend in blue-winged teal from 1986 to 1991 during Mid-winter Waterfowl Transects.	324
Figure IV.43. A. The number of blue-winged teal per count during Christmas Bird Counts. B. Trend in blue-winged teal from 1965 to 1989 during Christmas Bird Counts.	325
Figure IV.44. A. The number of scaup spp. per count during Christmas Bird Counts. B. Trend in scaup spp. from 1965 to 1989 during Christmas Bird Counts.	327
Figure IV.45. A. The number of ruddy ducks per count during Christmas Bird Counts. B. Trend in ruddy ducks from 1965 to 1989 during Christmas Bird Counts.	329
Figure IV.46. A. The number of buffleheads per count during Christmas Bird Counts. B. Trend in buffleheads from 1965 to 1989 during Christmas Bird Counts.	330
Figure IV.47. A. The number of wood ducks per count during Christmas Bird Counts. B. Trend in wood ducks from 1965 to 1989 during Christmas Bird Counts.	331
Figure IV.48. A. The number of ring-necked ducks per count during Christmas Bird Counts. B. Trend in ring-necked ducks from 1965 to 1989 during Christmas Bird Counts.	332
Figure IV.49. A. Individual counts and mean number of canvasbacks per count during Christmas Bird Counts. B. Individual counts and mean number of redheads per count during Christmas Bird Counts.	335

Figure IV.50. A. Mean numbers of red-winged blackbirds per stop during North American Breeding Bird Surveys. B. Mean numbers of boat-tailed and great-tailed grackles per stop during North American Breeding Bird Surveys.	336
Figure IV.51. A. Number of alligators per route and alligator nests per transect during Alligator Surveys. B. Trend in alligators per route from 1971 to 1984 during Alligator Surveys.	338
Figure IV.52. A. Total number of canvasbacks and ring-necked ducks during Mid-winter Waterfowl Cruise Counts. B. Total number of redheads and buffleheads during Mid-winter Waterfowl Cruise Counts. . . .	339
Figure IV.53 A. Total number of lesser scaups and American coots during Mid-winter Waterfowl Cruise Counts. B. Total number of ruddy ducks during Mid-winter Waterfowl Cruise Counts.	340
Figure V.1. Temporal distribution of phytoplankton studies in the Galveston Estuary from 1950 until 1990.	349
Figure V.2. Temporal distribution of zooplankton studies in the Galveston Estuary from 1950 until 1990.	349
Figure V.3. Temporal distribution of combined phyto- and zooplankton studies in the Galveston Estuary from 1950 until 1990.	350
Figure V.4. Variations in weekly means of chlorophyll <u>a</u> measurements in the Galveston Estuary during 1957-1959.	352
Figure V.5. Location of sampling stations in Cedar Bayou area and monthly chlorophyll <u>a</u> concentrations during 1972-1974.	353
Figure V.6. Locations of sampling stations in upper Galveston and Trinity Bays used in studies of Strong (1977), Krecji (1979) and Smith (1983).	354
Figure V.7. Monthly chlorophyll <u>a</u> concentrations from 10 stations in upper Galveston and Trinity Bays during 1976 (Strong 1977), 1977 (Krecji 1979) and 1978 (Smith 1983).	355
Figure V.8. Monthly primary productivity rates determined at ten stations in upper Galveston and Trinity Bays.	357
Figure VI.1. Marsh survey sites for macro-invertebrate infauna in the Galveston Estuary during 1987.	379

Figure VI.2. Abundances of total infauna and epifauna in sediment cores at marsh sites in the Galveston Estuary.	382
Figure VI.3. Abundances of annelids in sediment cores at marsh sites in the Galveston Estuary.	383
Figure VI.4. Abundances of peracarids in sediment cores at marsh sites in the Galveston Estuary.	384
Figure VI.5. Abundances of molluscs in sediment cores at marsh sites in the Galveston Estuary.	385
Figure VI.6. Abundances of miscellaneous infauna and epifauna in sediment cores at marsh sites in the Galveston Estuary.	387
Figure VI.7. Mean abundances of infauna in three habitats at a salt marsh in West Bay.	391
Figure VI.8. Mean abundances of annelids in three habitats at a salt marsh in West Bay.	392
Figure VI.9. Mean abundances of infaunal crustaceans in three habitats at a salt marsh in West Bay.	392
Figure VI.10. Mean abundances of infaunal mollusks in three habitats at a salt marsh in West Bay.	393
Figure VI.11. Mean abundances of other taxonomic groups in three habitats at a salt marsh in West Bay.	393
Figure VI.12. Mean abundances of <u>Strebosplio benedicti</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	394
Figure VI.13. Mean abundances of oligochaetes (Oligochaeta) in three habitats at a salt marsh in West Bay.	394
Figure VI.14. Mean abundances of <u>Capitella capitata</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	395
Figure VI.15. Mean abundances of <u>Mediomastus</u> spp. (Polychaeta) in three habitats at a salt marsh in West Bay.	395
Figure VI.16. Mean abundances of <u>Mediomastus californiensis</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	396
Figure VI.17. Mean abundances of <u>Mediomastus ambiseta</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	396

Figure VI.18. Mean abundances of <u>Polydora ligni</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	397
Figure VI.19. Mean abundances of <u>Nereis (Neanthes) succinea</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	397
Figure VI.20. Mean abundances of <u>Hobsonia florida</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	398
Figure VI.21. Mean abundances of <u>Heteromastus filiformis</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	398
Figure VI.22. Mean abundances of <u>Tharyx setigera</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	400
Figure VI.23. Mean abundances of <u>Melinna maculata</u> (Polychaeta) in three habitats at a salt marsh in West Bay.	400
Figure VI.24. Mean abundances of <u>Hargeria rapax</u> (Tanaidacea) in three habitats at a salt marsh in West Bay.	401
Figure VI.25. Mean abundances of <u>Corophium</u> sp. (Amphipoda) in three habitats at a salt marsh in West Bay.	401
Figure VI.26. Mean abundances of <u>Gammarus mucronatus</u> (Amphipoda) in three habitats at a salt marsh in West Bay.	402
Figure VI.27. Mean abundances of <u>Ampelisca abdita</u> (Amphipoda) in three habitats at a salt marsh in West Bay.	402
Figure VI.28. Mean abundances of <u>Edotea montosa</u> (Isopoda) in three habitats at a salt marsh in West Bay.	403
Figure VI.29. Mean abundances of <u>Grandidierella bonneroides</u> (Amphipoda) in three habitats at a salt marsh in West Bay.	403
Figure VI.30. Mean abundances of <u>Melita</u> cf. <u>nitida</u> (Amphipoda) in three habitats at a salt marsh in West Bay.	404
Figure VI.31. Mean abundances of <u>Orchestia</u> cf. <u>costericana</u> (Amphipoda) in three habitats at a salt marsh in West Bay.	404
Figure VI.32. Mean abundances of <u>Amygdalum papyrium</u> (Bivalvia) in three habitats at a salt marsh in West Bay.	405
Figure VI.33. Mean abundances of <u>Mulina lateralis</u> (Bivalvia) in three habitats at a salt marsh in West Bay.	405

Figure VI.34. Mean abundances of unidentified bivalves in three habitats at a salt marsh in West Bay.	406
Figure VI.35. Mean abundances of <u>Tagelus</u> sp. (Bivalvia) in three habitats at a salt marsh in West Bay.	406
Figure VII.1. Map of the Galveston Estuary showing areas referred to in the text.	418
Figure VII.2. Duration of monthly or semimonthly macrobenthic assemblage sampling programs at discrete localities within the Galveston Estuary.	420
Figure VII.3. Duration of monthly or semimonthly macrobenthic assemblage sampling programs at discrete localities within bay systems adjacent to the Galveston Estuary.	421
Figure VII.4. Trend of macrobenthic abundance in the Trinity Bay area in 1969.	427
Figure VII.5. Trends of macrobenthic abundances in the Trinity Bay area in 1970.	427
Figure VII.6. Trends of macrobenthic abundances in the Trinity Bay area in 1971.	428
Figure VII.7. Trend of macrobenthic abundance in the Trinity Bay area in 1972.	428
Figure VII.8. Trend of macrobenthic abundance in the Trinity Bay area in 1973.	429
Figure VII.9. Trend of macrobenthic abundance in the Houston Lighting and Power project, 1969-1973.	429
Figure VII.10. Trend of macrobenthic abundance in the Trinity Bay area in 1974.	430
Figure VII.11. Trend of macrobenthic abundance in the Trinity Bay area in 1975.	430
Figure VII.12. Trend of macrobenthic abundance in Clear Lake in 1970. . .	431
Figure VII.13. Trend of macrobenthic abundance in Clear Lake in 1971. . .	431
Figure VII.14. Trend of macrobenthic abundance in Eckert Bayou in 1976. . .	432

Figure VII.15. Trend of macrobenthic abundance in Eckert Bayou in 1977.	432
Figure VII.16. Trend of macrobenthic abundance in Eckert Bayou in 1978.	433
Figure VII.17. Trends of macrobenthic abundances in the West Bay area in 1980.	433
Figure VII.18. Trends of macrobenthic abundances in the West Bay area in 1981.	434
Figure VII.19. Comparison of quarterly macrobenthic abundances at five stations in the Galveston Estuary in 1971-72.	434
Figure VII.20. Trend of macrobenthic abundances in nearshore bottoms off Bolivar Roads in 1975-76.	437
Figure VII.21. Trend of macrobenthic abundance in Laguna del Oro, West Bay area, in 1980-81.	437
Figure VII.22. Trend of macrobenthic abundance in the Sea Rim State Park area in 1978-79.	438
Figure VII.23. Trend of macrobenthic abundance in Lavaca-Matagorda Bay in 1970-71.	438
Figure VII.24. Temporal trend of benthic abundance in Lavaca Bay in 1973-75.	441
Figure VII.25. Temporal trends of macrobenthic abundances in upper and lower San Antonio Bay in 1972-73.	441