TCEQ Interoffice Memorandum

То:	Lorinda Gardner, Regional Director Kent Waggoner, Air Section Manager David Ramirez, Border and Perimian Basin Area Director
From:	Tiffany Bredfeldt, Ph.D. TB Toxicology Division, Office of the Executive Director
Date:	October 19, 2012
Subject:	Health Effects Review of 2011 Ambient Air Network Monitoring Data in Region 6, El Paso

Conclusions

- Reported short-term concentrations of volatile organic compounds (VOCs) as detected by1hour automated gas chromatography (autoGC) would not be expected to cause adverse acute health effects, vegetation effects, or odorous conditions.
- Reported annual concentrations of VOCs, polycyclic aromatic hydrocarbons (PAHs), carbonyls, and metals reported as particulate matter with aerodynamic diameter of 2.5 microns or less (PM_{2.5}) and for one metal measured also in total suspended particulate matter (TSP) would also not be expected to cause long-term adverse human health or vegetation effects.

Background

This memorandum conveys the Toxicology Division's (TD) evaluation of ambient air sampling conducted at seven monitoring sites in Region 6-El Paso during 2011. TCEQ Region 6 monitoring site information is presented in Table 1 along with hyperlinks to detailed information regarding the monitoring sites and their maps. Lists 1-5, which can be found in Attachment A, display the target analytes for five monitoring sites. Three additional sites, Kern, Skyline Park, and Tillman, listed in Table 1 are the locations where lead TSP data were collected. The TD reviewed air monitoring summary results from 1-hour automated gas chromatography (autoGC) VOC samples, VOC canister samples collected on a 24-hour every sixth day schedule at Community Air Toxics Monitoring Network (CATMN) monitors, 24-hour metals samples (PM_{2.5} or TSP), 24-hour carbonyl samples, and 24-hour PAH or semivolatile organic compound (SVOC) samples.

The TCEQ Monitoring Division reported the data for all chemicals evaluated in this memorandum. Data discussed in this evaluation for all monitoring sites (84 VOCs from canister samples, 46 VOCs from autoGC, 17 carbonyls, 15 metals, and 16 PAHs), with the exception of the newest autoGC monitor at the Delta location, met the data completeness objective of 75 percent data return or at least 45 valid samples per year. Since 24-hour samples collected using the every sixth day schedule are designed to provide a representative long-term, ambient concentration for chemicals of concern, annual averages from all 24-hour samples were evaluated using appropriate long-term Air Monitoring Comparison Values (AMCVs) for the

Lorinda Gardner, et al. Page 2 October 19, 2012

potential to adversely impact long-term human health and vegetation effects. Thus, annual average concentrations of carbonyls, metals, PAHs, and VOCs (collected via canister sample), were compared to their respective long-term AMCVs. Reported annual average concentrations of lead (PM_{2.5} or TSP) were compared to the National Ambient Air Quality Standard (NAAQS). Annual average concentrations of VOCs collected via autoGC were also compared to their respective long-term AMCV. Hourly concentrations of VOCs collected by the autoGC were evaluated using the appropriate short-term AMCV. Additional information regarding the derivation and application of AMCVs is available <u>online</u>.

The El Paso Lower Valley monitor measures ambient concentrations of hydrogen sulfide (H_2S) and elevated levels of this chemical have been reported annually at this monitoring station since 2004. Further information regarding historical data collected at this monitoring station and subsequent evaluations of collected data are available from the Air Pollutant Watch List <u>website</u>.

City and Site Location	County	Monitor ID	Monitored Compounds
650 R.E. Thomason Loop (Ascarate Park SE)	El Paso	48-141-0055	VOCs ^a , Carbonyls, Metals (PM _{2.5})
800 S. San Marcial Street (El Paso Chamizal)	El Paso	48-141-0044	VOCs ^b , Carbonyls, Metals (PM _{2.5})
6700 Delta Drive (El Paso Delta)	El Paso	48-141-1011	VOCs ^b
700 San Francisco Ave (El Paso Sun Metro)	El Paso	48-141-0053	VOCs ^a , PAHs, Metals (PM _{2.5})
<u>301 East Robinson</u> (Kern)	El Paso	48-141-0033	Lead (TSP)
5050 A Yvette Drive (Skyline Park)	El Paso	48-141-0058	Lead (TSP)
<u>J. Harold Tillman HLT CT</u> 222 S Campbell S (Tillman)	El Paso	48-141-0002	Lead (TSP)
Clark & Cleveland Streets (Womble)	El Paso	48-141-0047	VOCs ^a

Table 1. Monitoring	Sites L	located in	TCEO	Region 6
	NICE DE			

^a24-hour canister only; ^bone-hour autoGC

Lorinda Gardner, et al. Page 3 October 19, 2012

Evaluation

VOCs

Hourly average concentrations of the 46 VOCs collected at the Chamizal and Delta autoGC monitoring site were below their respective short-term health-, odor-, and vegetation-based AMCVs. Thus, exposure to the reported hourly average concentrations would not be expected to cause adverse human health or welfare effects.

The 2011 annual average concentrations for all 84 VOCs collected as 24-hour canister samples at the Ascarate Park SE, Sun Metro and Womble monitoring sites were well below their respective long-term AMCVs. Because the Delta autoGC was activated in August of 2011, data collected at that site were not evaluated from a long-term perspective. Annual average concentrations for the 46 VOCs collected at the Chamizal autoGC monitoring site were also below their long-term AMCVs. Thus, adverse human health or vegetation effects would not be expected to occur as a result of long-term exposure to the reported levels of these chemicals at these monitoring sites.

Carbonyls

Reported annual average concentrations of the 17 carbonyls measured at the Ascarate Park SE monitoring site were below their respective long-term AMCVs and would not be expected to cause long-term adverse human health effects.

Metals

Reported annual average concentrations for all 14 metals (PM_{2.5}) measured at the Ascarate Park SE and Chamizal monitoring sites were below their respective AMCVs. In the case of lead (PM_{2.5} or TSP), reported annual average concentrations collected at the Ascarate Park SE, Chamizal, Kern, Skyline Park, and Tillman monitoring sites were below the lead NAAQS. Thus, none of the reported annual average concentrations for these15 metals (PM_{2.5} or TSP) would be expected to cause long term adverse health effects.

PAHs

The reported annual average concentrations for 15 of the 16 PAHs reported at the Sun Metro monitoring site in 2011 were well below their long-term AMCVs and would not be expected to cause long-term adverse human health effects. Phenanthrene was the only PAH with an annual average concentration (74.9 ng/m³) that exceeded the long-term AMCV (50 ng/m³). Figure 1 shows the annual average phenanthrene concentrations detected at the Sun Metro Monitoring Site from 1996 through 2011.

Lorinda Gardner, et al. Page 4 October 19, 2012




Figure 1. Phenanthrene Annual Average Concentrations at the Sun Metro Monitor for 16 Years. Note that bars are mean ± standard error. The 2012 data bar (*) represents only the first quarter phenanthrene concentrations.

Average annual concentrations of phenanthrene for 11 of the past 16 years, 1996, 1997, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, and 2011, respectively, have been greater than the long-term AMCV. The 16-year, which excludes 2012 data, average phenanthrene concentration of 66 ng/m³ is also greater than the long-term AMCV. The AMCV of 50 ng/m³ is a conservative interim value calculated from occupational exposure limits for a more toxic PAH, benzo(a) pyrene. Available literature investigating the toxicity of phenanthrene suggests that it is not carcinogenic to animals and is, therefore, significantly less toxic than the PAH benzo(a)pyrene, which is a known human carcinogen (Fertmann et al., 2002). The difference in toxicity and carcinogenicity appears to be due to metabolism. When phenanthrene enters human or animal bodies it is not metabolized to the same type of chemical intermediates that make benzo(a)pyrene carcinogenic (Hecht et al., 2010). Thus, based on toxicity data and the conservative nature of the long-term, health-based AMCV, we would not expect adverse effects on human health or welfare to occur as a result of exposure to the reported concentration of phenanthrene.

If you have any questions or comments regarding this evaluation, please feel free to contact me at (512) 239-1799 or <u>tiffany.bredfeldt@tceq.texas.gov</u>.

cc (via email):

Casso, Ruben – EPA Region 6, Dallas Prosperie, Susan – Department of State Health Services Lorinda Gardner, et al. Page 5 October 19, 2012

Attachment A

List 1. Target VOC Analytes in Canister Samples

1,1,2,2-Tetrachloroethane	Bromomethane	Methyl Chloroform (1,1,1-
1,1,2-Trichloroethane	Carbon Tetrachloride	Trichloroethane)
1,1-Dichloroethane	Chlorobenzene	Methylcyclohexane
1,1-Dichloroethylene	Chloroform	Methylcyclopentane
1,2,3-Trimethylbenzene	Chloromethane (Methyl	N-Butane
1,2,4-Trimethylbenzene	Chloride)	N-Decane
1,2-Dichloropropane	Cis 1,3-Dichloropropene	N-Heptane
1,3,5-Trimethylbenzene	Cis-2-Butene	N-Hexane
1,3-Butadiene	Cis-2-Hexene	N-Nonane
1-Butene	Cis-2-Pentene	N-Octane
1-Hexene+2-Methyl-1-Pentene	Cyclohexane	N-Pentane
1-Pentene	Cyclopentane	N-Propylbenzene
2,2,4-Trimethylpentane	Cyclopentene	N-Undecane
2,2-Dimethylbutane (Neohexane)	Dichlorodifluoromethane	O-Ethyltoluene
2,3,4-Trimethylpentane	Dichloromethane (Methylene	O-Xylene
2,3-Dimethylbutane	Chloride)	P-Diethylbenzene
2,3-Dimethylpentane	Ethane	P-Ethyltoluene
2,4-Dimethylpentane	Ethylbenzene	Propane
2-Chloropentane	Ethylene	Propylene
2-Methyl-2-Butene	Ethylene Dibromide (1,2-	Styrene
2-Methylheptane	Dibromoethane)	Tetrachloroethylene
2-Methylhexane	Ethylene Dichloride (1,2-	Toluene
2-Methylpentane (Isohexane)	Dichloroethane)	Trans-1-3-Dichloropropylene
3-Methyl-1-Butene	Isobutane	Trans-2-Butene
3-Methylheptane	Isopentane (2-Methylbutane)	Trans-2-Hexene
3-Methylhexane	Isoprene	Trans-2-Pentene
3-Methylpentane	Isopropylbenzene (Cumene)	Trichloroethylene
4-Methyl-1-Pentene	M-Diethylbenzene	Trichlorofluoromethane
Acetylene	M-Ethyltoluene	Vinyl Chloride
Benzene	M/P Xylene	-
		•

List 2. Target Carbonyl Analytes

Formaldehyde	o-Tolualdehyde
Heptaldehyde	Propanal - Propionaldehyde
Hexanaldehyde	p-Tolualdehyde
Isovaleraldehyde	Valeraldehyde
Methyl Ethyl Ketone	
(MEK)/Methacrolein	
m-Tolualdehyde	
	Heptaldehyde Hexanaldehyde Isovaleraldehyde Methyl Ethyl Ketone (MEK)/Methacrolein

List 3. Target Metal Analytes

Aluminum (PM_{2.5})

Antimony (PM 2.5)

Arsenic (PM_{2.5})

Lorinda Gardner, et al. Page 6 October 19, 2012

Barium (PM _{2.5})	Copper (PM _{2.5})	Nickel (PM _{2.5})
Cadmium (PM _{2.5})	Lead (PM _{2.5} or TSP)	Selenium (PM _{2.5})
Chromium (PM _{2.5})	Manganese(PM _{2.5})	Tin (PM _{2.5})
Cobalt (PM _{2.5})	Molybdenum (PM _{2.5})	Zinc $(PM_{2.5})$

List 4. Target PAH Analytes

Acenaphthene Acenaphthylene Anthracene Benzo (a) anthracene Benzo (a) pyrene Benzo (b) fluoranthene Molybdenum (PM_{2.5}) Benzo (ghi) perylene Benzo (k) fluoranthene Chrysene Dibenzo (a,h) anthracene

Fluoranthene

Fluorene

Indeno (1,2,3-cd) pyrene Naphthalene Phenanthrene Pyrene

List 5. Target VOC Analytes in AutoGC

1-Butene	Benzene	n-Heptane
1-Pentene	c-2-Butene	n-Hexane
1,2,3-Trimethylbenzene	c-2-Pentene	n-Nonane
1,2,4-Trimethylbenzene	Cyclohexane	n-Octane
1,3-Butadiene	Cyclopentane	n-Pentane
1,3,5-Trimethylbenzene	Ethane	n-Propylbenzene
2-Methylheptane	Ethyl Benzene	o-Xylene
2-Methylhexane	Ethylene	p-Xylene + m-Xylene
2,2-Dimethylbutane	Isobutane	Propane
2,2,4-Trimethylpentane	Isopentane	Propylene
2,3-Dimethylpentane	Isoprene	Styrene
2,3,4-Trimethylpentane	Isopropyl Benzene - Cumene	t-2-Butene
2,4-Dimethylpentane	Methylcyclohexane	t-2-Pentene
3-Methylheptane	Methylcyclopentane	Toluene
3-Methylhexane	n-Butane	
Acetylene	n-Decane	